

Stay Healthy This Flu Season!

It's that time of year when the crisp, cool air finally arrives, and along with it comes the influenza virus. Flu season typically begins in early fall and continues into winter, with peak flu season in January and February.



The flu is not the same as the common cold, but it is highly contagious and spreads in much the same way. The virus is contracted through particles in the air, on surfaces and by close contact with someone who is infected. It is an infection of the mouth, nose, throat and respiratory tract. The flu causes an array of symptoms: malaise, fever, chills, cough, sore throat, runny nose, congestion, headaches and muscle aches. Symptoms can last from days to weeks and may lead to complications such as ear infections, sinus infections, bronchitis and pneumonia.

IT'S TIME FOR YOUR FREE* FLU VACCINE!

DATE	TIME	LOCATION
10/09/2017	10:30-1:30	Central Office—Board Room
10/09/17	3:30-5:30	Central Office—Board Room

Everyone is at risk for contracting the flu, however, those at highest risk are young children (under the age of 5), the elderly (over the age of 65) and those that are immunosuppressed by certain medical conditions such as asthma, emphysema, diabetes, and heart disease. The severity of the flu varies from mild to severe and can lead to hospitalization and death. Even healthy individuals may suffer from complications of the flu.

There are many different strains of the flu virus and every year new vaccines are formulated to try to protect us from what are expected to be the top 3 or 4 strains of the virus. Flu vaccinations are our best protection against the flu virus. The vaccine reduces your risks of developing the flu and minimizes days of illness and lost productivity. Most importantly it can save lives.

The flu vaccine is usually available as early as August and recommended at any time during flu season, but vaccination is ideal before the end of October. Many people are afraid that the flu vaccine will cause the flu, however, the injectable forms of the flu virus are made with killed or inactivated forms of the flu virus and cannot cause an actual flu infection.

The most effective measure to reduce your risk of a flu infection this season is to get your flu vaccine. Talk to your doctor about the flu vaccine.

DID YOU KNOW?

Aetna members with the Whole Health plan (gold ID card) must get their flu vaccines from an in-network PCP, Urgent Care Center or on-site flu clinic provided by HealthTexas Medical Group



Article authored by **April Hain, M.D.**
 Board Certified in Family Medicine
 Top Doctors in America®
 Top Doctors in San Antonio®
[Additional Information on Dr. Hain](#)

Alamo Heights Clinic
 1248 Austin Hwy, Ste 214
 San Antonio, TX 78209
 Telephone: (210) 828-2531
 Fax: (210) 828-2532

*All vaccines will be billed to the member's insurance directly through HealthTexas Medical Group—Members must complete the registration form and bring their insurance card and ID as well.

**HealthTexas Medical Group is in-network and a preferred provider for the Aetna Whole Health (Baptist/HealthTexas Network)

Need a Primary Care Physician? Visit www.QPICSA.com or call (210) 920-9301

For more information on the flu season visit: <https://www.cdc.gov/flu/about/season/flu-season-2016-2017.htm>

Influenza (Flu) Vaccine (Inactivated or Recombinant): *What you need to know*

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Influenza (“flu”) is a contagious disease that spreads around the United States every year, usually between October and May.

Flu is caused by influenza viruses, and is spread mainly by coughing, sneezing, and close contact.

Anyone can get flu. Flu strikes suddenly and can last several days. Symptoms vary by age, but can include:

- fever/chills
- sore throat
- muscle aches
- fatigue
- cough
- headache
- runny or stuffy nose

Flu can also lead to pneumonia and blood infections, and cause diarrhea and seizures in children. If you have a medical condition, such as heart or lung disease, flu can make it worse.

Flu is more dangerous for some people. Infants and young children, people 65 years of age and older, pregnant women, and people with certain health conditions or a weakened immune system are at greatest risk.

Each year **thousands of people in the United States die from flu**, and many more are hospitalized.

Flu vaccine can:

- keep you from getting flu,
- make flu less severe if you do get it, and
- keep you from spreading flu to your family and other people.

2 Inactivated and recombinant flu vaccines

A dose of flu vaccine is recommended every flu season. Children 6 months through 8 years of age may need two doses during the same flu season. Everyone else needs only one dose each flu season.

Some inactivated flu vaccines contain a very small amount of a mercury-based preservative called thimerosal. Studies have not shown thimerosal in vaccines to be harmful, but flu vaccines that do not contain thimerosal are available.

There is no live flu virus in flu shots. **They cannot cause the flu.**

There are many flu viruses, and they are always changing. Each year a new flu vaccine is made to protect against three or four viruses that are likely to cause disease in the upcoming flu season. But even when the vaccine doesn’t exactly match these viruses, it may still provide some protection.

Flu vaccine cannot prevent:

- flu that is caused by a virus not covered by the vaccine, or
- illnesses that look like flu but are not.

It takes about 2 weeks for protection to develop after vaccination, and protection lasts through the flu season.

3 Some people should not get this vaccine

Tell the person who is giving you the vaccine:

- **If you have any severe, life-threatening allergies.**

If you ever had a life-threatening allergic reaction after a dose of flu vaccine, or have a severe allergy to any part of this vaccine, you may be advised not to get vaccinated. Most, but not all, types of flu vaccine contain a small amount of egg protein.

- **If you ever had Guillain-Barré Syndrome (also called GBS).**

Some people with a history of GBS should not get this vaccine. This should be discussed with your doctor.

- **If you are not feeling well.**

It is usually okay to get flu vaccine when you have a mild illness, but you might be asked to come back when you feel better.



4 Risks of a vaccine reaction

With any medicine, including vaccines, there is a chance of reactions. These are usually mild and go away on their own, but serious reactions are also possible.

Most people who get a flu shot do not have any problems with it.

Minor problems following a flu shot include:

- soreness, redness, or swelling where the shot was given
- hoarseness
- sore, red or itchy eyes
- cough
- fever
- aches
- headache
- itching
- fatigue

If these problems occur, they usually begin soon after the shot and last 1 or 2 days.

More serious problems following a flu shot can include the following:

- There may be a small increased risk of Guillain-Barré Syndrome (GBS) after inactivated flu vaccine. This risk has been estimated at 1 or 2 additional cases per million people vaccinated. This is much lower than the risk of severe complications from flu, which can be prevented by flu vaccine.
- Young children who get the flu shot along with pneumococcal vaccine (PCV13) and/or DTaP vaccine at the same time might be slightly more likely to have a seizure caused by fever. Ask your doctor for more information. Tell your doctor if a child who is getting flu vaccine has ever had a seizure.

Problems that could happen after any injected vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy, or have vision changes or ringing in the ears.
- Some people get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at about 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/

5 What if there is a serious reaction?

What should I look for?

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 and get the person to the nearest hospital. Otherwise, call your doctor.
- Reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor should file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not give medical advice.

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your healthcare provider. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/flu

Vaccine Information Statement
Inactivated Influenza Vaccine

08/07/2015

42 U.S.C. § 300aa-26

Office Use Only

